arXiv:1110.6759v2 [math.CO] 25 Mar 2012

π-FORMULAS WITH FREE PARAMETERS

^{A}CHUANAN WEI, ^{B}DIANXUAN GONG, ^{C} JIANBO LI

^{A}*Department of Information Technology*
Hainan Medical College, Haikou 571101, China
^{B}*College of Sciences*
Hebei Polytechnic University, Tangshan 063009, China
^{C}*Institute of Mathematical Sciences*
Xuzhou Normal University, Xuzhou 221116, China

Abstract. In terms of the hypergeometric method, we give the extensions of two known series for π. Further, other twenty-nine summation formulas for π, π^2 and $1/\pi$ with free parameters are also derived in the same way.

1. Introduction

For a complex number x and an integer n, define the shifted factorial by

$$(x)_n=\begin{cases}\prod_{i=0}^{n-1}(x+i), & \text{when} \quad n>0;\\ 1, & \text{when} \quad n=0;\\ \frac{1}{\prod_{j=n}^{-1}(x+j)}, & \text{when} \quad n<0.\end{cases}$$

Define Γ-function by Euler's integral:

$$\Gamma(x)=\int_0^{\infty}t^{x-1}e^{-t}dt \text{ with } Re(x)>0.$$

Then we have the following two relations:

$$\Gamma(x+n)=\Gamma(x)(x)_n, \quad \Gamma(x)\Gamma(1-x)=\frac{\pi}{\sin(\pi x)},$$

which will frequently be used without indication in this paper.

Following Bailey [4], define the hypergeometric series by

$${}_{1+r}F_s\left[\begin{matrix}a_0, & a_1, & \cdots, & a_r\\ & b_1, & \cdots, & b_s\end{matrix}\Big|\, z\right]=\sum_{k=0}^{\infty}\frac{(a_0)_k(a_1)_k\cdots(a_r)_k}{k!(b_1)_k\cdots(b_s)_k}z^k.$$

Then a simple ${}_2F_1$-series identity (cf. [30, Eq. (26)]) can be stated as

$${}_2F_1\left[\begin{matrix}1, & 1\\ & \frac{3}{2}\end{matrix}\Big|\, x^2\right]=\frac{\arcsin(x)}{x\sqrt{1-x^2}} \quad \text{where} \quad |x|<1.$$

2010 Mathematics Subject Classification: Primary 33C20 and Secondary 40A15, 65B10

Key words and phrases. Hypergeometric series; Summation formula for π; Ramanujan-type series for $1/\pi$.

Email addresses: weichuanan@yahoo.com.cn (C. Wei), gongdianxuan@yahoo.com.cn (D. Gong), ljianb66@gmail.com (J. Li).

Two beautiful series for π (cf. [30, Eq. (23) and (27)]) implied by it read as

$$\sum_{k=0}^{\infty}\frac{k!}{(2k+1)!!}=\frac{\pi}{2}, \tag{1}$$

$$\sum_{k=0}^{\infty}\frac{(k!)^2}{(2k+1)!}=\frac{2\pi}{3\sqrt{3}}. \tag{2}$$

Recall the ${}_7F_6$-series identity due to Chu [15, Eq. (5.1e)] and Dougall's ${}_5F_4$-series identity (cf. [4, p. 27]):

$$\begin{aligned}&{}_7F_6\left[\begin{matrix}a-\frac{1}{2}, & \frac{2a+2}{3}, & 2b-1, & 2c-1, & 2+2a-2b-2c, & a+s, & -s\\ & \frac{2a-1}{3}, & 1+a-b, & 1+a-c, & b+c-\frac{1}{2}, & 2a+2s, & -2s\end{matrix}\,\middle|\,1\right]\\ &=\frac{(\frac{1}{2}+a)_s(b)_s(c)_s(a-b-c+\frac{3}{2})_s}{(\frac{1}{2})_s(1+a-b)_s(1+a-c)_s(b+c-\frac{1}{2})_s},\end{aligned} \tag{3}$$

$$\begin{aligned}&{}_5F_4\left[\begin{matrix}a, & 1+\frac{a}{2}, & b, & c, & d\\ & \frac{a}{2}, & 1+a-b, & 1+a-c, & 1+a-d\end{matrix}\,\middle|\,1\right]\\ &=\frac{\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(1+a-d)\Gamma(1+a-b-c-d)}{\Gamma(1+a)\Gamma(1+a-b-c)\Gamma(1+a-b-d)\Gamma(1+a-c-d)}\end{aligned} \tag{4}$$

provided that $Re(1+a-b-c-d)>0$.

Recently, Chu [16, 17] and Liu [27, 28] deduced many surprising π-formulas from several known hypergeometric series identities. Thereinto, Chu [15] showed that (4) implies the Ramanujan-type series for $1/\pi$ with three free parameters:

$$\frac{2}{\pi}=\frac{(\frac{1}{2})_{m-n-p}}{(\frac{1}{2})_n(\frac{1}{2})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{1}{2})_{k+m}(\frac{1}{2})_{k+n}(\frac{1}{2})_{k+p}}{k!(k+m-n)!(k+m-p)!}(4k+2m+1) \tag{5}$$

where $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p\}\geq 0$ and Liu [28] showed that (4) implies the Ramanujan-type series for $1/\pi$ with four free parameters:

$$\begin{aligned}\frac{\sqrt{3}}{3\pi}&=\frac{(\frac{2}{3})_{m-n-p}(\frac{1}{3})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(m-n-p-q-1)!(\frac{1}{2})_n(\frac{1}{3})_p(\frac{2}{3})_q}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{2})_{k+m}(\frac{1}{2})_{k+n}(\frac{1}{3})_{k+p}(\frac{2}{3})_{k+q}}{k!(k+m-n)!(\frac{7}{6})_{k+m-p}(\frac{5}{6})_{k+m-q}}(1+2m+4k)\end{aligned} \tag{6}$$

where $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-n-p-q-1\}\geq 0$. More π-formulas can be found in the papers [1]-[3], [5]-[14], [18], [19]-[26] and [29]-[33].

Inspired by these work just mentioned, we shall explore further the relations of π-formulas and hypergeometric series. The structure of the paper is arranged as follows. Seven π-formulas with free parameters including the extensions of (1) and (2) will be derived from (3) in section 2. Twenty-four π-formulas with free parameters other than (5) and (6) will be deduced from (4) in section 3.

2. Summation formulas for π and π^2 with free parameters implied by Chu's ${}_7F_6$-series identity

Letting $s\to\infty$ for (3), we obtain the following equation:

$$\begin{aligned}&{}_5F_4\left[\begin{matrix}a-\frac{1}{2}, & \frac{2a+2}{3}, & 2b-1, & 2c-1, & 2+2a-2b-2c\\ & \frac{2a-1}{3}, & 1+a-b, & 1+a-c, & b+c-\frac{1}{2}\end{matrix}\,\middle|\,\frac{1}{4}\right]\\ &=\frac{\Gamma(\frac{1}{2})\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(b+c-\frac{1}{2})}{\Gamma(\frac{1}{2}+a)\Gamma(b)\Gamma(c)\Gamma(a-b-c+\frac{3}{2})}.\end{aligned} \tag{7}$$

Subsequently, one summation formula for π with two free parameters, five summation formulas for π with three free parameters and one summation formula for π^2 with three free parameters will respectively be derived from (7).

Choosing $b = 1 + m$, $c = 1 + n$ in (7) and then letting $a \to \infty$, we achieve the extension of (1).

Theorem 1. *For $m, n \in \mathbb{N}_0$, there holds the summation formula for π with two free parameters:*

$$\frac{\pi}{2^{m+n+1}} = \frac{m!n!}{(2m)!(2n)!} \sum_{k=0}^{\infty} \frac{(k+2m)!(k+2n)!}{k!(2k+2m+2n+1)!!}.$$

When $m = n = 0$, Theorem 1 reduces to (1) exactly. Other two examples of the same type are displayed as follows.

Example 1 ($m = 1$ and $n = 0$ in Theorem 1)**.**

$$\frac{\pi}{2} = \sum_{k=1}^{\infty} \frac{(k+1)!}{(2k+1)!!}.$$

Example 2 ($m = 2$ and $n = 0$ in Theorem 1)**.**

$$\frac{3\pi}{2} = \sum_{k=2}^{\infty} \frac{(k+2)!}{(2k+1)!!}.$$

Making $a = \frac{1}{2} + m$, $b = \frac{1}{4} + n$ and $c = \frac{3}{4} + p$ in (7), we attain the equation.

Theorem 2. *For $m, n, p \in \mathbb{Z}$ with $\min\{m, m-n-p\} \geq 0$, there holds the summation formula for π with three free parameters:*

$$\begin{aligned} \frac{\pi}{2} &= \frac{1}{(-\frac{1}{4})_n(\frac{1}{4})_p(\frac{1}{2})_{m-n-p}} \\ &\quad \times \sum_{k=0}^{\infty} \frac{(-\frac{1}{2})_{k+2n}(\frac{1}{2})_{k+2p}(k+m)!(k+2m-2n-2p)!}{(\frac{5}{4})_{k+m-n}(\frac{3}{4})_{k+m-p}(\frac{1}{2})_{k+n+p}\,k!} \frac{3k+2m}{k+m} \frac{1}{4^{k+m}}. \end{aligned}$$

Two examples from Theorem 2 are laid out as follows.

Example 3 ($m = n = 1$ and $p = 0$ in Theorem 2)**.**

$$\frac{\pi}{4} = \sum_{k=0}^{\infty} \frac{(2k)!}{(4k+3)!!}(3k+2).$$

Example 4 ($m = 2$ and $n = p = 1$ in Theorem 2)**.**

$$\frac{\pi}{4} = \sum_{k=1}^{\infty} \frac{(2k)!}{(4k+1)!!}(3k+1).$$

A beautiful result should be mentioned. Fixing $a = -\frac{3}{2}$ in the identity due to Chu [15, Equation (5.3 g)]:

$$\begin{aligned} & {}_6F_5\left[\begin{matrix} 7+4a, & -2-2a, & -\frac{5}{2}-2a, & \frac{2-4a}{5}, & -a+s, & -s \\ & \frac{4}{3}, & \frac{5}{3}, & -\frac{3+4a}{5}, & -2-4s, & -2-4a+4s \end{matrix} \;\middle|\; \frac{32}{27}\right] \\ = &\ \frac{(-\frac{1}{2}-a)_s(-\frac{1}{4}-a)_s(\frac{1}{4}-a)_s(\frac{5}{2}+a)_s}{(\frac{3}{4})_s(\frac{5}{4})_s(\frac{3}{2})_s(-\frac{3}{2}-2a)_s} \end{aligned}$$

and then letting $s \to \infty$, we obtain the surprising series for π:

$$\frac{\pi}{2} = \sum_{k=0}^{\infty} \frac{k!(2k)!}{(3k+2)!} \frac{5k+3}{2^k}.$$

Setting $a = \frac{1}{2} + m$, $b = \frac{1}{6} + n$ and $c = \frac{5}{6} + p$ in (7), we get the extension of (2).

Theorem 3. *For $m, n, p \in \mathbb{Z}$ with $\min\{m, m-n-p\} \geq 0$, there holds the summation formula for π with three free parameters:*

$$\begin{aligned}\frac{2\pi}{3\sqrt{3}} &= \frac{1}{(-\frac{1}{3})_n(\frac{1}{3})_p(\frac{1}{2})_{m-n-p}} \\ &\times \sum_{k=0}^{\infty} \frac{(-\frac{2}{3})_{k+2n}(\frac{2}{3})_{k+2p}(k+m)!(k+2m-2n-2p)!}{(\frac{4}{3})_{k+m-n}(\frac{2}{3})_{k+m-p}(\frac{1}{2})_{k+n+p}\,k!} \frac{3k+2m}{k+m} \frac{1}{4^{k+m}}.\end{aligned}$$

When $m = n = 1$ and $p = 0$, Theorem 3 reduces to (2) exactly. Other two examples of the same type are displayed as follows.

Example 5 ($m = 2$ and $n = p = 1$ in Theorem 3)**.**

$$\frac{2\pi}{3\sqrt{3}} = \sum_{k=1}^{\infty} \frac{(k!)^2}{(2k+1)!}(3k+2).$$

Example 6 ($m = 3$, $n = 2$ and $p = 1$ in Theorem 3)**.**

$$\frac{4\pi}{9\sqrt{3}} = \sum_{k=2}^{\infty} \frac{(k!)^2}{(2k+1)!}(3k+1)k.$$

Taking $a = \frac{1}{2} + m$, $b = \frac{1}{3} + n$ and $c = \frac{2}{3} + p$ in (7), we gain the equation.

Theorem 4. *For $m, n, p \in \mathbb{Z}$ with $\min\{m, m-n-p\} \geq 0$, there holds the summation formula for π with three free parameters:*

$$\begin{aligned}\frac{\pi}{\sqrt{3}} &= \frac{1}{(-\frac{1}{6})_n(\frac{1}{6})_p(\frac{1}{2})_{m-n-p}} \\ &\times \sum_{k=0}^{\infty} \frac{(-\frac{1}{3})_{k+2n}(\frac{1}{3})_{k+2p}(k+m)!(k+2m-2n-2p)!}{(\frac{7}{6})_{k+m-n}(\frac{5}{6})_{k+m-p}(\frac{1}{2})_{k+n+p}\,k!} \frac{3k+2m}{k+m} \frac{1}{4^{k+m}}.\end{aligned}$$

Two examples from Theorem 4 are laid out as follows.

Example 7 ($m = n = 1$ and $p = 0$ in Theorem 4)**.**

$$\frac{5\pi}{4\sqrt{3}} = \sum_{k=0}^{\infty} \frac{(1)_k(\frac{1}{3})_k(\frac{5}{3})_k}{(\frac{3}{2})_k(\frac{7}{6})_k(\frac{11}{6})_k} \frac{3k+2}{4^k}.$$

Example 8 ($m = 2$ and $n = p = 1$ in Theorem 4)**.**

$$\frac{\pi}{2\sqrt{3}} = \sum_{k=1}^{\infty} \frac{(1)_k(\frac{2}{3})_k(\frac{4}{3})_k}{(\frac{3}{2})_k(\frac{5}{6})_k(\frac{7}{6})_k} \frac{3k+1}{4^k}.$$

Choosing $a = \frac{1}{2} + m$, $b = \frac{5}{12} + n$ and $c = \frac{7}{12} + p$ in (7), we achieve the equation.

Theorem 5. *For $m, n, p \in \mathbb{Z}$ with $\min\{m, m-n-p\} \geq 0$, there holds the summation formula for π with three free parameters:*

$$\begin{aligned}\frac{\pi}{6(2-\sqrt{3})} &= \frac{1}{(-\frac{1}{12})_n(\frac{1}{12})_p(\frac{1}{2})_{m-n-p}} \\ &\times \sum_{k=0}^{\infty} \frac{(-\frac{1}{6})_{k+2n}(\frac{1}{6})_{k+2p}(k+m)!(k+2m-2n-2p)!}{(\frac{13}{12})_{k+m-n}(\frac{11}{12})_{k+m-p}(\frac{1}{2})_{k+n+p}\,k!} \frac{3k+2m}{k+m} \frac{1}{4^{k+m}}.\end{aligned}$$

Two examples from Theorem 5 are displayed as follows.

Example 9 ($m = n = 1$ and $p = 0$ in Theorem 5)**.**

$$\frac{11\pi}{60(2-\sqrt{3})} = \sum_{k=0}^{\infty} \frac{(1)_k(\frac{1}{6})_k(\frac{11}{6})_k}{(\frac{3}{2})_k(\frac{13}{12})_k(\frac{23}{12})_k} \frac{3k+2}{4^k}.$$

Example 10 ($m=2$ and $n=p=1$ in Theorem 5)**.**

$$\frac{\pi}{12(2-\sqrt{3})}=\sum_{k=1}^{\infty}\frac{(1)_k(\frac{5}{6})_k(\frac{7}{6})_k}{(\frac{3}{2})_k(\frac{11}{12})_k(\frac{13}{12})_k}\frac{3k+1}{4^k}.$$

Making $a=\frac{1}{2}+m$, $b=\frac{1}{12}+n$ and $c=\frac{11}{12}+p$ in (7), we attain the equation.

Theorem 6. *For $m,n,p\in\mathbb{Z}$ with $\min\{n,p,m-n-p\}\geq 0$, there holds the summation formula for π with three free parameters:*

$$\begin{aligned}\frac{5\pi}{6(2+\sqrt{3})} &= \frac{1}{(-\frac{5}{12})_n(\frac{5}{12})_p(\frac{1}{2})_{m-n-p}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(-\frac{5}{6})_{k+2n}(\frac{5}{6})_{k+2p}(k+m)!(k+2m-2n-2p)!}{(\frac{17}{12})_{k+m-n}(\frac{7}{12})_{k+m-p}(\frac{1}{2})_{k+n+p}\,k!}\frac{3k+2m}{k+m}\frac{1}{4^{k+m}}.\end{aligned}$$

Two examples from Theorem 6 are laid out as follows.

Example 11 ($m=n=1$ and $p=0$ in Theorem 6)**.**

$$\frac{35\pi}{12(2+\sqrt{3})}=\sum_{k=0}^{\infty}\frac{(1)_k(\frac{5}{6})_k(\frac{7}{6})_k}{(\frac{3}{2})_k(\frac{17}{12})_k(\frac{19}{12})_k}\frac{3k+2}{4^k}.$$

Example 12 ($m=2$ and $n=p=1$ in Theorem 5)**.**

$$\frac{5\pi}{12(2+\sqrt{3})}=\sum_{k=1}^{\infty}\frac{(1)_k(\frac{1}{6})_k(\frac{11}{6})_k}{(\frac{3}{2})_k(\frac{7}{12})_k(\frac{17}{12})_k}\frac{3k+1}{4^k}.$$

Setting $a=\frac{3}{2}+m$, $b=1+n$ and $c=1+p$ in (7), we get the equation.

Theorem 7. *For $m,n,p\in\mathbb{Z}$ with $\min\{n,p,m-n-p\}\geq 0$, there holds the summation formula for π^2 with three free parameters:*

$$\begin{aligned}\pi^2 &= \frac{1}{(\frac{1}{2})_n(\frac{1}{2})_p(\frac{1}{2})_{m-n-p}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(k+m)!(k+2n)!(k+2p)!(k+2m-2n-2p)!}{(\frac{3}{2})_{k+m-n}(\frac{3}{2})_{k+m-p}(\frac{3}{2})_{k+n+p}\,k!}\frac{3k+2m+2}{4^{k+m-1}}.\end{aligned}$$

Two examples from Theorem 7 are displayed as follows.

Example 13 ($m=n=p=0$ in Theorem 7)**.**

$$\frac{\pi^2}{4}=\sum_{k=0}^{\infty}\frac{(1)_k^3}{(\frac{3}{2})_k^3}\frac{3k+2}{4^k}.$$

Example 14 ($m=2$ and $n=p=1$ in Theorem 7)**.**

$$\frac{\pi^2}{12}=\sum_{k=2}^{\infty}\frac{(1)_k^3}{(\frac{1}{2})_k^2(\frac{3}{2})_k}\frac{k}{4^k}.$$

3. RAMANUJAN-TYPE SERIES FOR $1/\pi$ WITH FREE PARAMETERS IMPLIED BY DOUGALL'S ${}_5F_4$-SERIES IDENTITY

In this section, ten Ramanujan-type series for $1/\pi$ with three free parameters and fourteen Ramanujan-type series for $1/\pi$ with four free parameters other than (5) and (6) will respectively be deduced from (4).

Taking $a=\frac{1}{6}+m$, $b=\frac{1}{6}+n$, $c=\frac{1}{6}+p$ in (4) and then letting $d\to\infty$, we gain the equation.

Theorem 8. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p+1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3}{\pi}=\frac{(\frac{5}{6})_{m-n-p}}{(\frac{1}{6})_n(\frac{1}{6})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{1}{6})_{k+m}(\frac{1}{6})_{k+n}(\frac{1}{6})_{k+p}}{k!(k+m-n)!(k+m-p)!}(1+6m+12k).$$

Choosing $a=\frac{5}{6}+m$, $b=\frac{5}{6}+n$, $c=\frac{5}{6}+p$ in (4) and then letting $d\to\infty$, we achieve the equation.

Theorem 9. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3}{\pi}=\frac{(\frac{1}{6})_{m-n-p}}{(\frac{5}{6})_n(\frac{5}{6})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{5}{6})_{k+m}(\frac{5}{6})_{k+n}(\frac{5}{6})_{k+p}}{k!(k+m-n)!(k+m-p)!}(5+6m+12k).$$

Making $a=\frac{1}{4}+m$, $b=\frac{1}{4}+n$, $c=\frac{1}{4}+p$ in (4) and then letting $d\to\infty$, we attain the equation.

Theorem 10. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p+1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{2\sqrt{2}}{\pi}=\frac{(\frac{3}{4})_{m-n-p}}{(\frac{1}{4})_n(\frac{1}{4})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{1}{4})_{k+m}(\frac{1}{4})_{k+n}(\frac{1}{4})_{k+p}}{k!(k+m-n)!(k+m-p)!}(1+4m+8k).$$

Setting $a=\frac{3}{4}+m$, $b=\frac{3}{4}+n$, $c=\frac{3}{4}+p$ in (4) and then letting $d\to\infty$, we get the equation.

Theorem 11. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{2\sqrt{2}}{\pi}=\frac{(\frac{1}{4})_{m-n-p}}{(\frac{3}{4})_n(\frac{3}{4})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{3}{4})_{k+m}(\frac{3}{4})_{k+n}(\frac{3}{4})_{k+p}}{k!(k+m-n)!(k+m-p)!}(3+4m+8k).$$

Taking $a=\frac{1}{3}+m$, $b=\frac{1}{3}+n$, $c=\frac{1}{3}+p$ in (4) and then letting $d\to\infty$, we gain the equation.

Theorem 12. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3\sqrt{3}}{2\pi}=\frac{(\frac{2}{3})_{m-n-p}}{(\frac{1}{3})_n(\frac{1}{3})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{1}{3})_{k+m}(\frac{1}{3})_{k+n}(\frac{1}{3})_{k+p}}{k!(k+m-n)!(k+m-p)!}(1+3m+6k).$$

Choosing $a=\frac{2}{3}+m$, $b=\frac{2}{3}+n$, $c=\frac{2}{3}+p$ in (4) and then letting $d\to\infty$, we achieve the equation.

Theorem 13. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3\sqrt{3}}{2\pi}=\frac{(\frac{1}{3})_{m-n-p}}{(\frac{2}{3})_n(\frac{2}{3})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{2}{3})_{k+m}(\frac{2}{3})_{k+n}(\frac{2}{3})_{k+p}}{k!(k+m-n)!(k+m-p)!}(2+3m+6k).$$

Making $a=\frac{1}{12}+m$, $b=\frac{1}{12}+n$, $c=\frac{1}{12}+p$ in (4) and then letting $d\to\infty$, we attain the equation.

Theorem 14. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p+1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3(\sqrt{6}-\sqrt{2})}{\pi}=\frac{(\frac{11}{12})_{m-n-p}}{(\frac{1}{12})_n(\frac{1}{12})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{1}{12})_{k+m}(\frac{1}{12})_{k+n}(\frac{1}{12})_{k+p}}{k!(k+m-n)!(k+m-p)!}(1+12m+24k).$$

Setting $a=\frac{5}{12}+m$, $b=\frac{5}{12}+n$, $c=\frac{5}{12}+p$ in (4) and then letting $d\to\infty$, we get the equation.

Theorem 15. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3(\sqrt{6}+\sqrt{2})}{\pi}=\frac{(\frac{7}{12})_{m-n-p}}{(\frac{5}{12})_n(\frac{5}{12})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{5}{12})_{k+m}(\frac{5}{12})_{k+n}(\frac{5}{12})_{k+p}}{k!(k+m-n)!(k+m-p)!}(5+12m+24k).$$

Taking $a=\frac{7}{12}+m$, $b=\frac{7}{12}+n$, $c=\frac{7}{12}+p$ in (4) and then letting $d\to\infty$, we gain the equation.

Theorem 16. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3(\sqrt{6}+\sqrt{2})}{\pi}=\frac{(\frac{5}{12})_{m-n-p}}{(\frac{7}{12})_n(\frac{7}{12})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{7}{12})_{k+m}(\frac{7}{12})_{k+n}(\frac{7}{12})_{k+p}}{k!(k+m-n)!(k+m-p)!}(7+12m+24k).$$

Choosing $a=\frac{11}{12}+m$, $b=\frac{11}{12}+n$, $c=\frac{11}{12}+p$ in (4) and then letting $d\to\infty$, we achieve the equation.

Theorem 17. *For $m,n,p\in\mathbb{Z}$ with $\min\{m-n,m-p,m-2n-2p-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with three free parameters:*

$$\frac{3(\sqrt{6}-\sqrt{2})}{\pi}=\frac{(\frac{1}{12})_{m-n-p}}{(\frac{11}{12})_n(\frac{11}{12})_p}\sum_{k=0}^{\infty}(-1)^k\frac{(\frac{11}{12})_{k+m}(\frac{11}{12})_{k+n}(\frac{11}{12})_{k+p}}{k!(k+m-n)!(k+m-p)!}(11+12m+24k).$$

Making $a=\frac{1}{4}+m$, $b=\frac{1}{4}+n$, $c=\frac{1}{4}+p$ and $d=\frac{1}{2}+q$ in (4), we attain the equation.

Theorem 18. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-p,m-n-p-q\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{4}{\pi}&=\frac{(\frac{3}{4})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{1}{4})_{m-n-p-q}(\frac{1}{4})_n(\frac{1}{4})_p(\frac{1}{2})_q}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{4})_{k+m}(\frac{1}{4})_{k+n}(\frac{1}{4})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{3}{4})_{k+m-q}}(1+4m+8k).\end{aligned}$$

Setting $a=\frac{3}{4}+m$, $b=\frac{3}{4}+n$, $c=\frac{3}{4}+p$ and $d=\frac{1}{2}+q$ in (4), we get the equation.

Theorem 19. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-p,m-n-p-q-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{1}{\pi}&=\frac{(\frac{1}{4})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{3}{4})_{m-n-p-q-1}(\frac{3}{4})_n(\frac{3}{4})_p(\frac{1}{2})_q}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{3}{4})_{k+m}(\frac{3}{4})_{k+n}(\frac{3}{4})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{5}{4})_{k+m-q}}(3+4m+8k).\end{aligned}$$

Taking $a=\frac{1}{2}+m$, $b=\frac{1}{2}+n$, $c=\frac{1}{4}+p$ and $d=\frac{3}{4}+q$ in (4), we gain the equation.

Theorem 20. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-n-p-q-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{1}{2\pi}&=\frac{(\frac{3}{4})_{m-n-p}(\frac{1}{4})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(m-n-p-q-1)!(\frac{1}{2})_n(\frac{1}{4})_p(\frac{3}{4})_q}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{2})_{k+m}(\frac{1}{2})_{k+n}(\frac{1}{4})_{k+p}(\frac{3}{4})_{k+q}}{k!(k+m-n)!(\frac{5}{4})_{k+m-p}(\frac{3}{4})_{k+m-q}}(1+2m+4k).\end{aligned}$$

Choosing $a=\frac{1}{3}+m$, $b=\frac{1}{3}+n$, $c=\frac{1}{3}+p$ and $d=\frac{1}{2}+q$ in (4), we achieve the equation.

Theorem 21. *For $m, n, p, q \in \mathbb{Z}$ with $\min\{m-n, m-p, m-n-p-q\} \geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{3\sqrt{3}}{\pi} &= \frac{(\frac{2}{3})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{1}{6})_{m-n-p-q}(\frac{1}{3})_{n}(\frac{1}{3})_{p}(\frac{1}{2})_{q}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{3})_{k+m}(\frac{1}{3})_{k+n}(\frac{1}{3})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{5}{6})_{k+m-q}}(1+3m+6k).\end{aligned}$$

Making $a=\frac{2}{3}+m$, $b=\frac{2}{3}+n$, $c=\frac{2}{3}+p$ and $d=\frac{1}{2}+q$ in (4), we attain the equation.

Theorem 22. *For $m, n, p, q \in \mathbb{Z}$ with $\min\{m-n, m-p, m-n-p-q-1\} \geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{\sqrt{3}}{2\pi} &= \frac{(\frac{1}{3})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{5}{6})_{m-n-p-q-1}(\frac{2}{3})_{n}(\frac{2}{3})_{p}(\frac{1}{2})_{q}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(\frac{2}{3})_{k+m}(\frac{2}{3})_{k+n}(\frac{2}{3})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{7}{6})_{k+m-q}}(2+3m+6k).\end{aligned}$$

Setting $a=\frac{1}{6}+m$, $b=\frac{1}{6}+n$, $c=\frac{1}{6}+p$ and $d=\frac{1}{2}+q$ in (4), we get the equation.

Theorem 23. *For $m, n, p, q \in \mathbb{Z}$ with $\min\{m-n, m-p, m-n-p-q\} \geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{2\sqrt{3}}{\pi} &= \frac{(\frac{5}{6})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{1}{3})_{m-n-p-q}(\frac{1}{6})_{n}(\frac{1}{6})_{p}(\frac{1}{2})_{q}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{6})_{k+m}(\frac{1}{6})_{k+n}(\frac{1}{6})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{2}{3})_{k+m-q}}(1+6m+12k).\end{aligned}$$

Taking $a=\frac{5}{6}+m$, $b=\frac{5}{6}+n$, $c=\frac{5}{6}+p$ and $d=\frac{1}{2}+q$ in (4), we gain the equation.

Theorem 24. *For $m, n, p, q \in \mathbb{Z}$ with $\min\{m-n, m-p, m-n-p-q-1\} \geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{2\sqrt{3}}{3\pi} &= \frac{(\frac{1}{6})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{2}{3})_{m-n-p-q-1}(\frac{5}{6})_{n}(\frac{5}{6})_{p}(\frac{1}{2})_{q}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(\frac{5}{6})_{k+m}(\frac{5}{6})_{k+n}(\frac{5}{6})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{4}{3})_{k+m-q}}(5+6m+12k).\end{aligned}$$

Choosing $a=\frac{1}{2}+m$, $b=\frac{1}{2}+n$, $c=\frac{1}{6}+p$ and $d=\frac{5}{6}+q$ in (4), we achieve the equation.

Theorem 25. *For $m, n, p, q \in \mathbb{Z}$ with $\min\{m-n, m-n-p-q-1\} \geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{2\sqrt{3}}{9\pi} &= \frac{(\frac{5}{6})_{m-n-p}(\frac{1}{6})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(m-n-p-q-1)!(\frac{1}{2})_{n}(\frac{1}{6})_{p}(\frac{5}{6})_{q}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{2})_{k+m}(\frac{1}{2})_{k+n}(\frac{1}{6})_{k+p}(\frac{5}{6})_{k+q}}{k!(k+m-n)!(\frac{4}{3})_{k+m-p}(\frac{2}{3})_{k+m-q}}(1+2m+4k).\end{aligned}$$

Making $a=\frac{1}{12}+m$, $b=\frac{1}{12}+n$, $c=\frac{1}{12}+p$ and $d=\frac{1}{2}+q$ in (4), we attain the equation.

Theorem 26. *For $m, n, p, q \in \mathbb{Z}$ with $\min\{m-n, m-p, m-n-p-q\} \geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{12(2-\sqrt{3})}{\pi} &= \frac{(\frac{11}{12})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{5}{12})_{m-n-p-q}(\frac{1}{12})_{n}(\frac{1}{12})_{p}(\frac{1}{2})_{q}}\\ &\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{12})_{k+m}(\frac{1}{12})_{k+n}(\frac{1}{12})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{7}{12})_{k+m-q}}(1+12m+24k).\end{aligned}$$

Setting $a=\frac{5}{12}+m$, $b=\frac{5}{12}+n$, $c=\frac{5}{12}+p$ and $d=\frac{1}{2}+q$ in (4), we get the equation.

Theorem 27. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-p,m-n-p-q\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{12(2+\sqrt{3})}{\pi}&=\frac{(\frac{7}{12})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{1}{12})_{m-n-p-q}(\frac{5}{12})_{n}(\frac{5}{12})_{p}(\frac{1}{2})_{q}}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{5}{12})_{k+m}(\frac{5}{12})_{k+n}(\frac{5}{12})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{11}{12})_{k+m-q}}(5+12m+24k).\end{aligned}$$

Taking $a=\frac{7}{12}+m$, $b=\frac{7}{12}+n$, $c=\frac{7}{12}+p$ and $d=\frac{1}{2}+q$ in (4), we gain the equation.

Theorem 28. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-p,m-n-p-q-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{2+\sqrt{3}}{\pi}&=\frac{(\frac{5}{12})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{11}{12})_{m-n-p-q-1}(\frac{7}{12})_{n}(\frac{7}{12})_{p}(\frac{1}{2})_{q}}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{7}{12})_{k+m}(\frac{7}{12})_{k+n}(\frac{7}{12})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{13}{12})_{k+m-q}}(7+12m+24k).\end{aligned}$$

Choosing $a=\frac{11}{12}+m$, $b=\frac{11}{12}+n$, $c=\frac{11}{12}+p$ and $d=\frac{1}{2}+q$ in (4), we achieve the equation.

Theorem 29. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-p,m-n-p-q-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{5(2-\sqrt{3})}{\pi}&=\frac{(\frac{1}{12})_{m-n-p}(\frac{1}{2})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(\frac{7}{12})_{m-n-p-q-1}(\frac{11}{12})_{n}(\frac{11}{12})_{p}(\frac{1}{2})_{q}}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{11}{12})_{k+m}(\frac{11}{12})_{k+n}(\frac{11}{12})_{k+p}(\frac{1}{2})_{k+q}}{k!(k+m-n)!(k+m-p)!(\frac{17}{12})_{k+m-q}}(11+12m+24k).\end{aligned}$$

Making $a=\frac{1}{2}+m$, $b=\frac{1}{2}+n$, $c=\frac{1}{12}+p$ and $d=\frac{11}{12}+q$ in (4), we attain the equation.

Theorem 30. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-n-p-q-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{5(2-\sqrt{3})}{6\pi}&=\frac{(\frac{11}{12})_{m-n-p}(\frac{1}{12})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(m-n-p-q-1)!(\frac{1}{2})_{n}(\frac{1}{12})_{p}(\frac{11}{12})_{q}}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{2})_{k+m}(\frac{1}{2})_{k+n}(\frac{1}{12})_{k+p}(\frac{11}{12})_{k+q}}{k!(k+m-n)!(\frac{17}{12})_{k+m-p}(\frac{7}{12})_{k+m-q}}(1+2m+4k).\end{aligned}$$

Setting $a=\frac{1}{2}+m$, $b=\frac{1}{2}+n$, $c=\frac{5}{12}+p$ and $d=\frac{7}{12}+q$ in (4), we get the equation.

Theorem 31. *For $m,n,p,q\in\mathbb{Z}$ with $\min\{m-n,m-n-p-q-1\}\geq 0$, there holds the Ramanujan-type series for $1/\pi$ with four free parameters:*

$$\begin{aligned}\frac{2+\sqrt{3}}{6\pi}&=\frac{(\frac{7}{12})_{m-n-p}(\frac{5}{12})_{m-n-q}(\frac{1}{2})_{m-p-q}}{(m-n-p-q-1)!(\frac{1}{2})_{n}(\frac{5}{12})_{p}(\frac{7}{12})_{q}}\\&\quad\times\sum_{k=0}^{\infty}\frac{(\frac{1}{2})_{k+m}(\frac{1}{2})_{k+n}(\frac{5}{12})_{k+p}(\frac{7}{12})_{k+q}}{k!(k+m-n)!(\frac{13}{12})_{k+m-p}(\frac{11}{12})_{k+m-q}}(1+2m+4k).\end{aligned}$$

Remark: With the change of the parameters, Theorems 8-31 can produce numerous concrete Ramanujan-type series for $1/\pi$. We shall not lay them out one by one here.

REFERENCES

[1] V. Adamchik, S. Wagon, A simple formula for π, Amer. Math. Monthly 104 (1997), 852-855.

[2] D.H. Bailey, J.M. Borwein, Experimantal mathematics: examples, methods and implications, Notices Amer. Math. Soc. 52 (2005), 502-514.

[3] D.H. Bailey, P.B. Borwein, S. Plouffe, On the rapid computation of various polylogarithmic constants, Math. Comp. 66 (1997), 903-913.

[4] W.N. Bailey, Generalized Hypergeometric Series, Cambridge University Press, Cambridge, 1935.

[5] N.D. Baruah, B.C. Berndt, Ramanujan's series for $1/\pi$ arising from his cubic and quartic theories of elliptic functions, J. Math. Anal. Appl. 341 (2008), 357-371.

[6] N.D. Baruah, B.C. Berndt, Eisenstein series and Ramanujan-type series for $1/\pi$, Ramanujan J. (2009), DOI 10.1007/s11139-008-9155-8.

[7] N.D. Baruah, B.C. Berndt, Ramanujan's Eisenstein series and new hypergeometric-like series for $1/\pi^2$, J. Approx. Theory 160 (2009), 135-153.

[8] N.D. Baruah, B.C. Berndt, H.H. Chan, Ramanujan's series for $1/\pi$: a survey, Amer. Math. Monthly 116 (2009), 567-587.

[9] J.M. Borwein, P.B. Borwein, π and the AGM: A Study in Analytic Number Theory and Computotional Complexity, Wiley, New York, 1987.

[10] J.M. Borwein, P.B. Borwein, Class number three Ramanujan type series for $1/\pi$, J. Comput. Appl. Math. 46 (1993), 281-290.

[11] H.C. Chan, More formulas for π, Amer. Math. Monthly 113 (2006), 452-455.

[12] H.H. Chan, S.H. Chan, Z.G. Liu, Domb's numbers and Ramanujan-Sato type series for $1/\pi$, Adv. Math. 186 (2004), 396-410.

[13] H.H. Chan, W.C. Liaw, Cubic modular equations and new Ramanujan-type series for $1/\pi$, Pacific J. Math. 192 (2000), 219-238.

[14] H.H. Chan, W.C. Liaw, V. Tan, Ramanujan's class invariant λ_n and a new class of series for $1/\pi$, J. London Math. Soc. 64 (2001), 93-106.

[15] W. Chu, Inversion techniques and combinatorial identities: A unified treatment for the ${}_7F_6$-series identities, Collect. Math. 45 (1994), 13-43.

[16] W. Chu, π-formulas implied by Dougall's summation theorem for ${}_5F_4$-series, Ramanujan J. 26 (2011), 251-255.

[17] W. Chu, Dougall's bilateral ${}_2H_2$-series and Ramanujan-like π-formulae, Math. Comp. 80 (2011), 2223-2251.

[18] J.W.L. Glaisher, On series for $1/\pi$ and $1/\pi^2$, Q.J. Math. 37 (1905), 173-198.

[19] B. Gourévitch, J. Guillera, Construction of binomial sums for π and polylogarithmic constants inspired by BBP formulas, Appl. Math. E-Notes 7 (2007), 237-246.

[20] J. Guillera, About a new kind of Ramanujan-type series, Exper. Math. 12 (2003), 507-510.

[21] J. Guillera, A new method to obtain series for $1/\pi$ and $1/\pi^2$, Exper. Math. 15 (2006), 83-89.

[22] J. Guillera, A class of conjectured series representations for $1/\pi$, Exper. Math. 15 (2006), 409-414.

[23] J. Guillera, Generators of some Ramanujan formulas, Ramanujan J. 11 (2006), 41-48.

[24] J. Guillera, History of the formulas and algorithms for π(Spanish), Gac. R. Soc. Mat. Esp. 10 (2007), 159-178.

[25] J. Guillera, Hypergeometric identities for 10 extended Ramanujan-type series, Ramanujan J. 15 (2008), 219-234.

[26] P. Levrie, Using Fourier-Legendre expansions to derive series for $1/\pi$ and $1/\pi^2$, Ramanujan J. 22 (2010), 221-230.

[27] Z. Liu, Gauss summation and Ramanujan type series for $1/\pi$, Int. J. Number Theory 8 (2012) 289-297.

[28] Z. Liu, A summation formula and Ramanujan type series, J. Math. Anal. Appl. 389 (2012) 1059-1065.

[29] S. Ramanujan, Modular equations and approximations to π, Quart. J. Pure Appl. Math. 45 (1914), 350-372.

[30] W.E. Weisstein, Pi Formulas, MathWorld-A Wolfram Web Resource, http:// mathworld.wolfram.com/PiFormulas.html.

[31] D. Zheng, Multisection method and further formulae for π, Indian J. Pure Ap. Math. 139 (2008), 137-156.

[32] W. Zudilin, More Ramanujan-type formulas for $1/\pi^2$(Russian), Usp. Mat. Nauk. 62 (2007), 211-212.

[33] W. Zudilin, Ramanujan-type formulas for $1/\pi$: A second wind?, arXiv: math. NT/0712. 1332V2.